



Carvacrol in oregano: the key compound and the role of thymol in the plant

Dr. A. Colonnese¹, R. Panzironi²

¹ Nutrition Biologist, Latina, Italy

² Independent Researcher, Latina, Italy

1. ABSTRACT

When we talk about oregano, we often think of its aroma, but its chemistry is much richer. Two substances, carvacrol and thymol, explain a large part of the plant's profile and the differences between oregano varieties. Carvacrol is among the most studied compounds and is a phenolic monoterpene (an aromatic molecule produced by plants). Its chemical form can change how it acts in biological systems. Thymol completes this picture and can modify the effects of carvacrol. To understand the data, one must distinguish between the spice, the essential oil, and concentrated extracts. These products differ in composition and potency. In this article, we will look at these molecules, their variety, and the phytocomplex (the set of substances in the plant). We will also see how much the form of consumption matters. In summary, here are the key points we will explore:

- Carvacrol and thymol define a large part of oregano's chemical profile.
- Spice, essential oil, and extracts are different products.
- Composition changes with species, harvest, and processing method.
- The phytocomplex matters more than a single molecule taken in isolation.
- Caution and a clear context are needed to interpret the data correctly.

VIDEO ARTICLE



ARTICLE INFORMATION

Received: October 22, 2026

Revised: October 22, 2026

Accepted: October 22, 2026

Published: October 2026

DOI 10.5281/zenodo.placeholder

KEYWORDS

carvacrolo origano
timolo origano
olio essenziale origano composizione
composti fenolici origano
principi attivi origano
carvacrolo e timolo

MISSION

LIFE Science Hub – Journal of Science was founded with the aim of making scientific knowledge accessible without giving up methodological rigour, quality of sources and authoritativeness of content. The journal promotes the dissemination of knowledge in the health, nutrition, medical and longevity sciences, building a bridge between the world of research and society.

Through articles, reviews, in-depth analyses and expert contributions, the Journal champions evidence-based medicine and science, fostering a correct interpretation of scientific data and countering misinformation. Every piece of content is developed with the aim of translating the complexity of research into understandable, useful and applicable information, while keeping intact the principles of accuracy, transparency and scientific independence.

Free from political, industrial and commercial influence, the journal stands as an authoritative and independent editorial space, dedicated to promoting scientific culture, professional development and the growth of health awareness among the public.



2. CARVACROL AND THYMOL: WHAT THEY ARE AND WHY THEY CHANGE THE IDENTITY OF OREGANO

In oregano, carvacrol and thymol define much of the plant's aroma and biological profile. These are phenolic monoterpenes (small aromatic plant molecules), which are therefore fragrant and highly reactive compounds. These substances can influence the scent and the way the phytocomplex—which includes all the substances naturally present in the plant—behaves. For this reason, oregano should not be viewed as a spice that is always the same. Discussing oregano carvacrol therefore means starting with two key molecules and their ratio. This balance matters more than just the presence of the plant itself. In fact, it can change how oregano is perceived and studied as a source of active ingredients. In the next step, this same pair of compounds must be evaluated in different forms: spice, extract, and essential oil. Before distinguishing between the various products, it is useful to establish a few essential points:

- Carvacrol and thymol are phenolic compounds found in oregano.
- Their presence influences the aroma and sensory identity.
- The composition varies between different types of oregano and among various processing methods.
- The phytocomplex allows for the evaluation of multiple active ingredients together.

To understand the chemical identity of oregano, one must first know its guiding molecules. The following sections describe the two most typical compounds and their roles. Their quantities, in fact, are not fixed and can vary significantly.

2.1 What are carvacrol and thymol in oregano

These two key compounds are phenolic monoterpenes (small aromatic molecules of plant origin). Their structure significantly influences the chemical profile of oregano. Within the plant, they contribute to its scent and the perception of its intensity. For this reason, they are among the most characteristic active ingredients of the *Origanum* genus. However, their significance also depends on their ratio to other substances in the phytocomplex,

which provides a useful criterion for describing an individual chemical profile with greater precision. Studies and technical reports indicate that these quantities vary based on species, botanical origin, and processing [1][2]. A 2011 analysis examined an oregano essential oil. Carvacrol was at 36.46%, thymol at 29.74%, and p-cymene at 24.31% [3]. A 2018 review then described the former as a monoterpene phenolic typical of the plant. The same source notes that human evidence remains incomplete [4]. These data therefore describe specific profiles, not every oregano in absolute terms.

2.2 Why the chemical profile of oregano is not always the same

The profile of oregano changes because there is no single composition that applies to all plants, all seasons, and all processing methods. Oregano essential oil composition and the spice's content depend on botanical and environmental factors, as well as harvesting, drying, and storage. This is why two products sold under the same name may show different levels of carvacrol, thymol, and other minor constituents. A 2024 review on Lamiaceae confirmed that these two compounds are among the dominant ones in most matrices, but not with the same distribution everywhere [5]. The useful takeaway is practical: variability does not disprove the role of oregano's phenolic compounds, but it requires that every extract be interpreted within its own context. The limitation, here too, is that summary data cannot replace the analysis of an individual sample.

2.3 Oregano phytocomplex: why it matters more than the individual compound

The phytocomplex includes all the substances naturally present in a plant. Together, these molecules form its final profile. In oregano, this concept avoids reducing everything to a single compound. This holds true even when carvacrol and thymol are present in higher quantities. Aroma, stability, and part of the biological activity may depend on the ratios between multiple substances, rather than an isolated molecule. A 2016 review studied the ethanolic extract of *Origanum vulgare*. The extract showed cytotoxic (harmful to cells), antioxidants, and antibacterial activity. The authors linked these effects primarily to



the two lead compounds [6]. The active ingredients of oregano should therefore be viewed as a system, not as separate parts. However, tests on laboratory extracts do not directly describe the spice or oils on the market. Bioavailability (the portion used by the body) and the extraction method can, in fact, change the picture.

3. FROM SPICE TO ESSENTIAL OIL: HOW COMPOSITION AND CONCENTRATION CHANGE

When moving from the kitchen to concentrated products, the quantity, form, and use change. The composition of oregano essential oil does not match that of the spice. In concentrates, the phenolic compounds of oregano are present in higher doses. Among these are phenolic monoterpenes, small aromatic molecules that have been widely studied. This distinction prevents incorrect comparisons between dietary use and concentrated preparations. The portion of active ingredients absorbable by the body may also change. Consequently, the value of the results obtained in experiments changes. An effect observed in an extract or an oil does not automatically apply to the spice. The following paragraphs explain how to interpret extracts and why the various forms are not interchangeable [7][3]. In the spice, however, the phytocomplex (the set of substances in the plant) remains diluted in the food. Therefore, carvacrol and thymol can have different proportions in the various forms. Before reading scientific data, it is necessary to distinguish between form, extraction method, and declared composition:

- The culinary spice is a complex and diluted mixture.
- Essential oil concentrates volatile compounds and phenols.
- Extracts vary based on the method used to obtain them.
- Labels help to understand the declared composition of the product.

Before examining the studies, it is useful to distinguish between the forms of oregano cited in the research. The same plant, in fact, can provide products that are very different in terms of composition, dose, and use.

3.1 Spice, essential oil, and extracts: three non-equivalent products

Culinary oregano, essential oil, and commercial extracts are not equivalent products. The spice contains limited amounts of the most studied compounds. Essential oil, on the other hand, is a concentrated part obtained through extraction (the separation of certain substances). Extracts can vary in purity, dosage, and the ratio between molecules. Standardization—the guaranteed presence of a specific quantity—can also vary. These parameters determine how much of the studied substance is actually present in the sample analyzed in the laboratory. This difference matters because the effect of carvacrol and thymol depends on the actual dose and the form used. A 2024 systematic review evaluated 38 articles on the two compounds. The studies reported antibiofilm (against layers of adherent microbes) and antimicrobial activity. There were also 68 synergistic combinations, with enhanced effects, or additive ones, with summed effects [8]. However, this data refers to preparations used in experiments, not the household spice. Distinguishing between these forms therefore avoids overly broad conclusions.

3.2 How to read oregano extract labels

Labels may indicate the standardization of the extract and the amount of its main compounds. Not all products, however, report the same data or present it with equal clarity. A complete label distinguishes a simple reference to the plant from a concentrate with a declared composition. This information helps to understand which product is being used and in what dose. Without these details, the name oregano alone does not accurately describe the content. This data is essential for comparing different products without immediately assuming they are equivalent. In 2020, a thymol chemotype essential oil (a variant rich in this substance) was described as potentially suitable for selective herbicidal use on monocotyledonous species [9]. The result shows that the chemotype can significantly change the oil's properties. However, this concerns a possible herbicidal use, not a health effect. The chemical label therefore guides interpretation, but it does not make oil and spice comparable.



3.3 When spice and essential oil cannot be compared

Comparing a spice and an essential oil as equivalent products can lead to incorrect conclusions. Spices are used in small doses and consumed along with other foods. Their compounds are therefore much more diluted compared to those in a concentrated product. An essential oil, on the other hand, collects volatile substances and is intended for different uses. Even with the same plant, the dosage, bioavailability (the portion absorbable by the body), and method of use change. A comparison therefore requires considering concentration, the context of use, and the specific product studied together. Effects observed in a laboratory or with concentrated products do not automatically apply to cooking. In 2021, a systematic review examined the anti-tumor effects of carvacrol and thymol. The authors reiterated that these are preclinical data, obtained before human studies, and should be interpreted with caution [10]. These premises help explain why one of the two compounds receives so much attention in research.

4. WHAT MAKES CARVACROL INTERESTING FROM A BIOLOGICAL PERSPECTIVE

Carvacrol is a phenolic monoterpene (a small plant molecule with a phenolic group). In study models, it can act on multiple biological targets. Tests evaluate both its action on microbes and its effects on membranes and oxidative stress. The latter occurs when too many reactive substances form within the cell. The composition of oregano essential oil also matters, as it can vary significantly. The phytocomplex (the set of substances in the plant) changes based on the raw material and the extraction method. The ratio between the phenolic compounds in oregano also varies. Carvacrol is therefore an important part of a larger system. Its role is clearly visible in tests, but caution is required when discussing food or human use. These data explain why the compound is useful for studying the biological activity of oregano. However, they do not directly demonstrate a clinical benefit. To better understand the topic, it is helpful to start with three questions that will guide the next steps:

- Why carvacrol has become the most studied compound in oregano.
- How carvacrol can act on cell membranes.
- Why results change when moving from the molecule to the plant.

Carvacrol helps explain why oregano generates so much interest in biological research. The following sections show its role, how it acts, and the limitations of comparing it to the whole plant.

4.1 Why carvacrol is the most studied molecule in oregano

Carvacrol is widely studied because it shows various easily measurable effects in tests. It is a phenolic monoterpene (a small plant molecule with a phenolic group). Its structure allows it to act on multiple parts of the cell, rather than having a single effect. Furthermore, certain chemotypes (plant types with different chemical compositions) contain higher amounts of it. This is why researchers often use it to carefully study the active ingredients of oregano. This choice allows for linking the molecule's structure to the effects observed in various tests. A 2018 review summarized these data and the biological role of the compound. However, it clarified that human trials (studies conducted on people) are still lacking [4]. Its interest therefore remains primarily biological and related to testing, rather than clinical treatment. The strength of the results depends on the model used. Tests help to understand the mechanisms, but they are not sufficient to define a therapeutic use in humans. One of the first effects studied concerns cell membranes.

4.2 Carvacrol and cell membranes: the first level of action

Carvacrol is also being studied for how it interacts with cell membranes. The molecule is lipophilic (it binds well to fats) and can enter their lipid bilayer. It can thus partially change the shape, fluidity, and order of the membrane. In some microbes, this can reduce their stability and alter exchanges with the outside environment. This is a physical and chemical effect observed in tests, not proof of a clinical effect. An in vitro study (conducted in a laboratory) examined *Pseudomonas aeruginosa* and *Staphylococcus aureus*. Researchers tested oregano essential oil and its main



components. They then measured the minimum amount capable of stopping bacterial growth [11]. The result shows that the effect on the membrane can produce measurable biological activity. However, the evidence does not indicate the effect in humans. It also does not clarify what happens within the phytocomplex, where other substances may increase or reduce the action.

4.3 Why comparing an isolated molecule to the plant is complicated

Studying carvacrol on its own is useful, but it is not enough to understand what really happens when using oregano. In the plant, in fact, the compound coexists with other constituents that can modify its stability, absorption, and overall biological response. These constituents do not remain inert: they can strengthen an effect or attenuate it, depending on the quantities involved. For this reason, the comparison between an isolated molecule and a plant matrix is always delicate. A pure substance helps to clarify the mechanism with precision, isolating one variable at a time. The spice or oil, on the other hand, describes a more complex and less linear system, made up of many molecules acting together. In 2025, a review also described synergistic antimicrobial activities between carvacrol and thymol against bacteria and *Candida* [12]. Synergy refers to a combined effect that is stronger than the sum of the individual effects. This data reinforces the idea that oregano should be viewed as a system of molecules, not as a single isolated active ingredient. However, the limitation remains interpretive: these data support a relevant biological role, but they do not allow for the direct transfer of results from the isolated molecule to the use of the whole plant. Caution is therefore needed when moving from the laboratory to real-world consumption.

5. HOW TO INTERPRET THE ROLE OF THYMOL IN OREGANO

Context is necessary to understand the role of thymol in oregano and its weight within the mixture. Thymol contributes to the plant's properties, but it does not explain its value on its own. It must be interpreted alongside the other active ingredients, because the final profile depends on the entire phytocomplex. This method avoids overly simplistic interpretations of the

composition of oregano essential oil. It also explains why different samples may have aromas and functions that are not entirely identical. Thymol remains important, but it describes the plant better when compared with other phenolic monoterpenes. Furthermore, harvesting, processing, and the method used to extract the substances also matter. To interpret the data correctly, it is useful to follow four steps:

- Evaluate thymol as part of the total composition.
- Study the ratio with carvacrol in the phytocomplex.
- Distinguish the composition of the oil from the biological response.
- Consider how extraction changes the proportion of the compounds.

Thymol is a phenolic monoterpene (an aromatic terpene molecule with a phenolic group). It should be viewed as part of the chemical profile of oregano, not as an independent substance.

5.1 The role of thymol in the biological profile of oregano

Thymol contributes to the oregano profile as one of the main phenolic compounds, but its significance must always be considered in relation to the rest of the matrix. In practice, it describes a portion of the aromatic signature and potential biological activity, without encompassing the entire function of the plant. The reference review [7] shows that the concentration of thymol can range from trace amounts to over 60% of the essential oil, depending on the subspecies and geographical origin. This wide range explains why two oregano samples can have very different sensory and biological profiles. The collected data also indicate that thymol shows antimicrobial activity in vitro, but the observed potency varies based on the total composition of the oil. The limitation of this interpretation is clear: the work synthesizes literature data and does not measure a single sample, so while it provides a good general overview, it does not allow for attributing a universal or identical effect to thymol across every botanical origin.

5.2 Carvacrol and thymol: how the ratio between varieties changes

The ratio between carvacrol and thymol is useful because it describes how the composition of oregano changes from one chemotype to another. In some

cases, carvacrol prevails, while in others, thymol is more prominent, and this difference affects both the scent and the overall biological profile. The studies collected in the review [7] document chemotypes with a carvacrol/thymol ratio greater than 10 to 1 in some variants of *Origanum vulgare*, while in others, thymol dominates with a ratio of less than 1 to 2. These imbalances are not minor details: they influence the antioxidant capacity measured with the DPPH assay and modify the spectrum of action against Gram-positive and Gram-negative bacteria. For this reason, talking about only one of the two molecules risks oversimplifying a variable chemical reality. The limitation remains that of descriptive and preclinical research: they indicate real differences, but they are not enough on their own to predict the practical effect of a specific product.

5.3 Why thymol should not be interpreted alone

Isolating thymol can be useful for studying its

properties, but within the plant, its significance emerges primarily in conjunction with other constituents. In the oregano phytocomplex, the interactions between molecules, their relative abundance, and the way the matrix is obtained are what truly matter. The review [7] cites evidence of synergy between thymol and carvacrol, where the mixture of the two compounds shows antimicrobial activity superior to that of the individual components tested at the same concentration. This combined effect is lost when evaluating a molecule isolated from its natural context. This also applies when comparing the spice, extract, and essential oil, because the bioavailability of the components changes significantly between the different forms. The data show a genuine interest in this combined action, but they remain almost entirely preclinical, with inconsistent results across models. The most solid takeaway is therefore interpretative: thymol should not be viewed as an absolute indicator, but as part of a broader chemical network.

6. QUADRO COMPARATIVO E SINTESI DELLE EVIDENZE

This summary synthesizes the previously cited literature. It connects the composition, the variety of products, and the activities observed in laboratory tests.

Sintesi delle evidenze scientifiche

Publication year	Area studied	What the studies indicate
2001	antimicrobial mechanism of action	in vitro effects were observed on <i>Pseudomonas aeruginosa</i> and <i>Staphylococcus aureus</i> with essential oil, thymol, and carvacrol [11]
2011	chemical composition of an oregano chemotype	in the analyzed sample, 36.46% carvacrol, 29.74% thymol, and 24.31% p-cymene were measured [3]
2016	ethanolic extract of <i>Origanum vulgare</i> and main components	evidence attributes much of the antioxidant and antibacterial activity to carvacrol and thymol [6]
2018	biological interest in carvacrol	studies indicate broad interest in testing, but clinical data on humans remain incomplete [4]
2020	guideline on oregano essential oil	evidence indicates that the essential oil is a defined mixture, different from the spice and rich in carvacrol and thymol [7]
2023	variability of composition based on species and cultivation	studies suggest that carvacrol and thymol levels vary significantly with botanical origin and growing conditions [1]



2024	antibacterial and antibiofilm activity of carvacrol and thymol	38 articles were evaluated, with mean minimum inhibitory concentrations of 475.46 and 279.26 µg/mL and 68 synergistic or additive combinations [8]
2025	antimicrobial synergy between carvacrol and thymol	data indicate a combined action against bacteria and Candida, but these are still derived from preclinical studies [12]

7. SINTESI EDITORIALE E CONCLUSIONI

EDITORIAL INTERPRETATION OF THE EVIDENCE

In the studies I reviewed for this article, I found one point on which the literature converges. Carvacrol and thymol are reliable markers for studying oregano. I observed that the composition changes significantly between species, chemotypes, and products. Furthermore, the observed effects are highly dependent on the dose and the matrix. In laboratory tests, these compounds show activity against certain microbes and other biological actions.

The main limitation I encountered is the gap between laboratory tests and real-world use. Much of the data comes from reviews or in vitro trials (outside of a living organism). I noted that samples, extraction methods, and analysis techniques vary between studies. Moreover, the spice, essential oil, and extracts are not equivalent products. To interpret the results, I therefore believe it is essential to clearly define the form, dose, and context.

FINAL CRITICAL NOTE

The overview I have outlined describes the chemistry of oregano well, but provides fewer answers regarding its practical use. The studies I have examined confirm the central role of carvacrol and thymol and the great variety in composition. However, many comparable data on humans are still missing. Therefore, a gap remains between what we know about the chemistry and what happens in real-world use. Comparing the studies also proved difficult for me. Indeed, the species, extraction methods, dosages, and test models vary. The most accurate interpretation I can offer remains cautious: the topic has a solid foundation, but more data are still needed.

CONCLUSIONS AND EDITORIAL SUMMARY

From the studies gathered here, I have come to the conviction that carvacrol and thymol help in studying oregano with care, as they define much of its chemical identity. They also explain many differences between samples, chemotypes (chemical variants of the same species), and products. I have observed that these compounds do not act alone. The matrix, the phytocomplex, and the actual dose also matter. For this reason, the spice, essential oil, and extracts are not equivalent. The tests help me understand the potential biological role of oregano. However, laboratory data do not apply directly to everyday life. Studying aromatic plants remains useful in nutrition because it links quality, natural variety, and the careful use of food. Oregano therefore offers a good model for evaluating the bioactive compounds (substances that act within biological systems) of plants.

Aggiornato al 22/10/2026. Questo contenuto riflette una sintesi divulgativa delle evidenze scientifiche al momento della pubblicazione e non sostituisce il parere medico.

8. TRANSPARENCY AND SCIENTIFIC RESPONSIBILITY

Artificial Intelligence was used solely as a tool for editorial and documentary support. All content, interpretations, bibliographic citations and conclusions have been verified, reviewed and approved by the author, who assumes full scientific responsibility for them.

The manuscript was also subjected to independent peer review prior to publication.

To ensure the greatest transparency, the complete history of the article, including the dates of submission, revision, acceptance and publication, as well as information on the peer review process, is available on the dedicated Article History page.



9. RIFERIMENTI BIBLIOGRAFICI

- [1] **Limited Scope Technical Report: Flavors - Handling**

2023

USDA

DOI:

<https://www.ams.usda.gov/sites/default/files/media/LimitedScopeTechnicalReportFlavorsHandling2023.pdf>

- [2] **Oregano Essential Oil as an Antimicrobial and Antioxidant Additive in Food Products**

Rodriguez-Garcia et al., 2016

Critical Reviews in Food Science and Nutrition

DOI: <https://doi.org/10.1080/10408398.2013.800832>

- [3] **Oregano: Chemical Analysis and Evaluation of Its Antimalarial, Antioxidant, and Cytotoxic Activities**

Babili et al., 2011

Journal of Food Science

DOI: <https://doi.org/10.1111/j.1750-3841.2011.02109.x>

- [4] **Carvacrol and human health: A comprehensive review**

Sharifi Rad et al., 2018

Phytotherapy Research

DOI: <https://doi.org/10.1002/ptr.6103>

- [5] **Chemical and Bioactive Evaluation of Essential Oils from Edible and Aromatic Mediterranean Lamiaceae Plants**

Spréa et al., 2024

Molecules

DOI: <https://doi.org/10.3390/molecules29122827>

- [6] **Antioxidant, Antibacterial, and Cytotoxic Activities of the Ethanolic *Origanum vulgare***

Extract and Its Major Constituents

Coccimiglio et al., 2016

Oxidative Medicine and Cellular Longevity

DOI: <https://doi.org/10.1155/2016/1404505>

- [7] **Safety and efficacy of an essential oil of *Origanum vulgare* ssp. *hirtum* (Link) leetsw. for all poultry species**

2020

EFSA

DOI: <https://www.efsa.europa.eu/en/efsajournal/pub/6195>

- [8] **Thymol and carvacrol against *Klebsiella*: anti-bacterial, anti-biofilm, and synergistic activities—a systematic review**

Farhadi et al., 2024

Frontiers in Pharmacology

DOI: <https://doi.org/10.3389/fphar.2024.1487083>

- [9] **Thymol Chemotype *Origanum vulgare* L. Essential Oil as a Potential Selective Bio-Based Herbicide on Monocot Plant Species**

Gruľová et al., 2020

Molecules

DOI: <https://doi.org/10.3390/molecules25030595>

- [10] **Antitumor Effects of Carvacrol and Thymol: A Systematic Review**

Sampaio et al., 2021

Frontiers in Pharmacology

DOI: <https://doi.org/10.3389/fphar.2021.702487>

- [11] **A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol**

Lambert et al., 2001

Journal of Applied Microbiology

DOI: <https://doi.org/10.1046/j.1365-2672.2001.01428.x>

- [12] **Carvacrol and Thymol, a Synergistic Antimicrobial Activity Against Bacterial and *Candida***

Species

Radocchia et al., 2025

MicrobiologyOpen

DOI: <https://doi.org/10.1002/mb03.70089>



- [13] **Biological and Pharmacological Activities of Carvacrol and Carvacrol Bearing Essential Oils**

Can Baser et al., 2008

Current Pharmaceutical Design

DOI: <https://doi.org/10.2174/138161208786404227>

- [14] **Mexican Oregano (*Lippia graveolens* Kunth) as Source of Bioactive Compounds: A Review**

Bautista-Hernández et al., 2021

Molecules

DOI: <https://doi.org/10.3390/molecules26175156>

- [15] **"In vivo" and "in vitro" antimicrobial activity of *Origanum vulgare* essential oil and its two phenolic compounds on clinical isolates of *Candida* spp.**

Stringaro et al., 2023

Archives of Microbiology

DOI: <https://doi.org/10.1007/s00203-022-03355-1>