LIFE Science Hub
JOURNAL OF SCIENCE
EXPANDING KNOWLEDGE ABOUT NUTRITION AND HEALTH

ORIGINAL RESEARCH
Volume 1, Issue 4, November 2026
ISSN 2845-7501

Insulin resistance: what it is, how it develops, and why it impacts daily metabolism

Dott. A. Conte[1]

1 Nutritional Biologist, Italy

1. ABSTRACT

Insulin resistance is a key topic for metabolism. It indicates that tissues respond less effectively to the insulin signal. This hormone helps the body use glucose, which is blood sugar. It also helps keep glycemia stable. If cells respond less, the body must produce more insulin to compensate. Consequently, it regulates energy less precisely. This phenomenon is not just a laboratory finding. It involves muscle, liver, and adipose tissue, which manage glucose and fats in different ways. Understanding insulin resistance helps separate the biological mechanism from its clinical effects. It also allows for a better evaluation of metabolic studies. In particular, this article addresses:

- What a reduced response to insulin really indicates
- Which tissues help control glucose
- Why the insulin signal can lose effectiveness
- Which metabolic signals can guide the assessment
- Why this framework helps to better interpret studies

ONLINE ARTICLE

ARTICLE INFORMATION

Published: November 19, 2026
Updated: November 19, 2026
Reviewed: November 19, 2026

DOI 10.5281/zenodo.24365450

KEYWORDS

insulino-resistenza
sensibilità all'insulina
metabolismo glucidico
glucosio nel sangue
tessuto adiposo
fegato

MISSION

LIFE Science Hub – Journal of Science was founded with the aim of making scientific knowledge accessible without giving up methodological rigour, quality of sources and authoritativeness of content. The journal promotes the dissemination of knowledge in the health, nutrition, medical and longevity sciences, building a bridge between the world of research and society.

Through articles, reviews, in-depth analyses and expert contributions, the Journal champions evidence-based medicine and science, fostering a correct interpretation of scientific data and countering misinformation. Every piece of content is developed with the aim of translating the complexity of research into understandable, useful and applicable information, while keeping intact the principles of accuracy, transparency and scientific independence.

Free from political, industrial and commercial influence, the journal stands as an authoritative and independent editorial space, dedicated to promoting scientific culture, professional development and the growth of health awareness among the public.

2. WHAT INSULIN RESISTANCE REALLY INDICATES

We speak of insulin resistance when tissues respond less to the insulin signal. This hormone helps glucose enter cells, where it is used. Its metabolic signal therefore becomes weaker.Insulin resistance does not always coincide with a single diagnosis. It indicates a process that changes over time and can precede more evident disorders. This distinction helps to interpret metabolism without confusing the cause with the result. Let us now see which tissues are involved and why the insulin signal may lose strength. The body may produce more insulin or struggle more to manage glucose after meals. This gradually changes the balance of glycemia, or blood glucose. To better understand the problem, it is useful to follow a few interconnected steps:

- The response to insulin drops, but it can vary between tissues in the same body.
- Initially, the body can compensate and keep blood glucose stable.
- The effort increases when the system manages the glucose consumed during meals.
- To evaluate the clinical picture, multiple signals are needed, because a single value is not enough.

Before examining the tissues, it is necessary to understand why insulin acts less effectively and how metabolism changes.

2.1 What insulin resistance really means

Insulin resistance indicates a reduced effectiveness of insulin in transmitting its message to the tissues. The hormone's signal arrives, but the tissues respond more weakly. It is not a single diagnosis, because it can appear in various conditions and with varying intensity. Often, the body produces more hormone to achieve the same effect on glucose and energy use. It is therefore a mechanism that can change over time, rather than a fixed clinical label. The HOMA-IR is an index based on fasting glycemia and insulin levels. A study on its average values in healthy subjects [1] shows that reference ranges vary between individuals. Even with normal glycemia, the body can be under stress and compensating without clear symptoms. However, a single data point does not describe the entire picture, because compensation can change over time. The same person can, in fact, go through various phases of compensation or loss of efficiency.

2.2 Difference between insulin resistance, hyperinsulinemia, and high glycemia

Insulin resistance, hyperinsulinemia, and high blood sugar refer to three different phenomena, so they are not synonyms. The first indicates that tissues respond more weakly to the insulin signal. The second indicates more hormone in the blood, often because the body produces it to compensate. The third indicates a measurable increase in blood glucose, also called glycemia. This distinction separates the mechanism from its outcome and helps in interpreting tests in the early stages. A review on insulin resistance in adipose tissue [2] shows that the defect can appear before a clear increase in glycemia. In the meantime, the hormone can increase and help the body still maintain this balance. This compensation can hide the problem for a long time and make it more difficult to notice. However, no single value distinguishes a short-term adaptation from a more stable disorder. It is therefore necessary to evaluate all available data together to better understand the picture.

3. WHICH TISSUES RESPOND LESS WELL TO INSULIN

The ability to respond well to insulin indicates how effectively tissues react to this hormone's signal. The response varies across different parts of the body because each performs specific tasks. In insulin resistance, muscle, liver, and adipose tissue no longer manage the metabolic signal effectively.Each tissue impacts the final balance differently. Muscle uses the majority of glucose after meals. The liver primarily regulates fasting glycemia. Adipose tissue, on the other hand, helps manage energy and hormonal signals. Let us now examine these three points one by one, keeping their roles distinct. The problem, therefore, does not concern only blood glucose. It involves multiple tissues and takes different forms in each. To understand the overall picture, it is useful to distinguish the roles that change when the signal loses strength:

- Muscle absorbs less glucose after meals.
- The liver reduces its internal glucose production less effectively.

- Adipose tissue sends less balanced metabolic signals.
- The crosstalk between tissues alters overall glycemic control.

The tissues involved in insulin resistance do not all respond poorly in the same way. Every part of the body has a specific metabolic task. Therefore, the effects emerge at different times of the day.

3.1 The role of muscle in glucose uptake

Skeletal muscle responds to insulin and allows glucose to enter cells after meals. This is the mechanism that enables the body to effectively use energy from meals. When this response decreases, the body uses this fuel less efficiently. Consequently, glucose remains in the blood for longer. It is not just about "high blood sugar," but about a less effective use of glucose. In fact, the muscle does not absorb or burn glucose as it should. Since muscle makes up a large mass, even a slight decline can affect overall glycemia control. A small, widespread change is enough to raise average blood values. Reviews describe various changes that reduce glucose uptake [3]. These changes involve the internal cellular pathways that respond to insulin. This data helps to understand the central role of muscle in insulin resistance. However, this tissue does not act alone. Physical activity, available energy, and communication with the liver and adipose tissue change the picture. These factors can improve or worsen muscle response over time. Its clinical significance therefore stems from its central position, not from an isolated defect. In practice, it is the primary consumer of post-prandial (after-meal) glucose [3].

3.2 How the liver contributes to fasting glycemia

The liver regulates fasting glycemia and produces glucose between meals. Insulin acts as a brake on this production. When the liver's response to insulin decreases, the brake loses its strength. The organ then continues to release glucose, even when the body does not need it. This process raises blood glucose during fasting, when levels should remain more stable. For this reason, the reduced liver response to insulin has a significant impact on values observed between meals. A review shows that signals from adipose tissue can promote hepatic insulin resistance. The same study therefore links adipose signals to two aspects of the liver's metabolic function. This condition changes glucose production and fat storage in the organ [4]. Hepatic control, however, does not depend solely on insulin. Available energy and other metabolic signals also play a role. The picture therefore reflects an altered balance, not a simple on-off switch. In practice, the liver primarily affects glycemia measured while fasting [4].

3.3 Adipose tissue and global metabolic balance

Adipose tissue is not just an energy store: it is an organ that sends hormonal and metabolic signals to the rest of the body. When its function changes, the regulation of energy balance becomes less efficient, and insulin struggles more to coordinate the various tissues. This happens because the hormones produced by fat travel to the liver and muscles. There, they modulate how these organs use glucose. If the signal is unbalanced, other tissues are affected as well. In this sense, adipose tissue contributes to the balance of glycemia not only by storing fat, but also by modulating low-grade inflammation and communication with the liver and muscles. Mild but continuous inflammation can, in fact, interfere with the insulin signal in nearby tissues. The interpretive limit is that it is not just the amount of adipose tissue that counts: its distribution, biological activity, and inflammatory state are also relevant. Two people with the same weight can therefore have different metabolic responses. For this reason, its role is not reduced to body weight, but to the ability to maintain an orderly metabolic dialogue.

4. WHY THE INSULIN SIGNAL WEAKENS

The insulin signal loses strength when it encounters repeated obstacles. The body must then produce more insulin to achieve the same effect. In insulin resistance , there is no single cause. Multiple factors reduce the cells' ability to respond to insulin. Consequently, the body struggles more to control blood glucose. It is a slow change, not a sudden failure.These factors do not act alone, but reinforce each other. In this way, they reduce glycemic homeostasis (the balance of glucose in the blood). We will first see how excess fat hinders the response to insulin. Then, we will understand how mild inflammation and daily habits can prolong this

state. To better understand this process, it is useful to distinguish the main mechanisms:

- Excess fat changes the function of adipose tissue and the liver.
- Low-grade inflammation alters the signals within cells.
- Lifestyle and sleep change the body's response every day.
- Individual predisposition modifies the level of personal risk.

Insulin signaling loses strength because multiple mechanisms act on the same control pathway. The following paragraphs explain how fats, inflammation, and habits change the body's response.

4.1 Fat accumulation and resistance to insulin response

When adipose tissue accumulates too much energy, its communication with the rest of the body changes. It is not just the amount of fat that matters, but also where it is located and how it is stored. The tissue must, in fact, store it without impairing glucose utilization. If adipose storage functions poorly, the liver receives more substances than it can manage. Consequently, the response to insulin becomes less precise. Reviews link visceral fat and excess lipids (fats) to insulin resistance [5][6]. The connection grows when adipose tissue does not effectively retain fatty acids and the signals that promote inflammation. However, studies do not indicate a threshold that applies to everyone. The same amount of fat can have different effects depending on its location and duration. The state of the liver also matters. The problem therefore arises when the storage, transport, and use of energy lose their balance.

4.2 Low-grade inflammation and metabolic communication

Low-grade inflammation keeps the immune system slightly active, but for a long time. It does not cause the clear signs typical of acute inflammation. In this state, the signals from cells that regulate glucose become less orderly. The response to insulin also loses effectiveness. It is not a single substance at work. Instead, what matters is the environment created around cells, vessels, and tissues. This widespread disturbance makes the metabolic message passing between different tissues less clear. Studies indicate that signs of inflammation and insulin resistance can reinforce each other [7][8]. This creates a cycle that maintains metabolic damage. However, this relationship varies from person to person. Inflammation can worsen the condition or stem from the metabolic problem. In some cases, it can play both roles. For this reason, the data describe a reciprocal link, without establishing an identical sequence for everyone. It should therefore be viewed as a factor that amplifies the process, not as its sole cause. Daily habits can also promote or limit this state.

4.3 Lifestyle, sleep, and individual predisposition

The insulin response depends on the tissues, but also on how we live our daily lives. Physical activity, diet, sleep, and personal predisposition influence blood glucose control. If these factors remain unfavorable for a long time, the body uses energy with less flexibility. The ability to respond to insulin therefore tends to decrease. These elements act together and can stabilize an inefficient metabolic response. Available evidence associates more movement and less time spent sitting with better metabolic control. Conversely, a sedentary lifestyle promotes a less effective response to insulin over time. The effect, however, is not the same for everyone. Sleep, family history, and initial status can significantly change an individual's reaction. Data therefore help to estimate risk, but they do not predict the individual case with certainty. Therefore, the observed associations describe general trends, not a predetermined destiny for everyone. Lifestyle alone does not create the problem, but it can significantly change its course.

5. HOW TO RECOGNIZE THE ASSOCIATED METABOLIC PROFILE

The metabolic profile linked to insulin resistance does not emerge from a single value. It is necessary to evaluate multiple signals that indicate a less effective use of sugars. This process is called glucose metabolism (use of sugars). The body can maintain relative glycemic homeostasis (blood sugar balance) through compensation. Meanwhile, blood glucose levels may fluctuate more. The liver, muscles, and adipose tissue must also work harder. For this reason, medical consultations and tests must go beyond a

single data point.These elements are part of the same metabolic dysfunction (metabolic alteration). They are, therefore, not isolated pieces of evidence. The following sections explain which signals may appear and why the problem may remain barely visible at first.This is where the limitations of self-interpretation emerge. Without the proper context, some data may appear normal even if the metabolic signal is already altered. To recognize this profile, some key clues are useful:

- Glycemia may become less stable after meals.
- Insulin in the blood may increase to compensate for the problem.
- Blood fats may show an altered balance.
- Adipose tissue may make the insulin signal less effective.
- The liver may play a more prominent role in the overall picture.

The picture becomes clearer when you connect the signals, rather than observing them one at a time. The following steps describe which metabolic clues may appear. They also explain when it is not advisable to stop at the first impression.

5.1 Metabolic signals that may accompany insulin resistance

The signals most often accompanying insulin resistance concern the body's ability to manage blood glucose and distribute energy in an orderly manner. An increase in glycemia after meals, higher circulating insulin, and alterations in the lipid profile may appear together or at different times. In parallel, adipose tissue and the liver participate in this less stable balance. Their workload increases to compensate for a less responsive reaction to insulin. Thus, the weight of insulin resistance on daily metabolism becomes more evident. Data collected in meta-analyses show that these signals tend to present as a set. They do not always follow the same order, but they share a common basis. For this reason, they should be read as parts of a single picture, not as separate events. The central point is that these are not diagnostic signals taken individually, but clues consistent with a less robust balance. The limitation is that these studies are primarily syntheses: they help define the pattern, but they do not transform a single data point into an autonomous diagnosis.

5.2 Why the disorder can remain silent for a long time

The situation may remain barely visible because, in the initial stages, the body compensates for the reduced ability to respond to insulin by increasing the workload of certain tissues and maintaining relative stability in glucose metabolism. This means that blood glucose may still appear within expected limits, despite an increasing biological cost. The body works harder to achieve the same apparent result. The tissues involved increase their activity, but their margin decreases over time. As long as the compensation holds, routine tests may not show anything unusual. This explains why the problem can go unnoticed for long periods. The silence does not indicate the absence of a problem: it indicates, rather, that the altered metabolic signaling is temporarily masked by adaptation mechanisms. The limitation of the available data is that they describe the mechanism well, but do not allow for precise prediction of when the condition will become evident in an individual. For this reason, clinical silence should not be mistaken for harmlessness.

5.3 When the condition requires a clinical evaluation

Clinical evaluation becomes important when metabolic signals are not episodic, but persist or are associated with other risk factors. In that case, the goal is not to immediately label the person, but to understand whether the combination of blood glucose, insulin, and lipid profile describes a true metabolic dysfunction, that is, an alteration in how the metabolism functions. Individual values must be placed within the person's history. It is necessary to observe how they change over time and how they relate to one another. Only in this way can a transient moment be distinguished from a more stable trend. Professional interpretation is primarily useful for organizing data and distinguishing a temporary adaptation from a condition that requires attention. Available summaries indicate that measuring how well the body responds to insulin may require more data and more context, because no single parameter is enough to capture the complexity of the clinical picture [9]. The limitation remains practical: tests help provide guidance, but the correct interpretation

depends on the set of information and the timeframe in which it is observed.

6. WHY THIS CLINICAL PICTURE MATTERS WHEN DISCUSSING TURMERIC

When discussing turmeric, one cannot take an effect on metabolism for granted. It is necessary to understand the biological framework in which it is being studied. Insulin resistance is part of a network that regulates the response to insulin and blood glucose. Many tissues act together within this network.In research, therefore, turmeric is studied in the presence of altered glycemic homeostasis (the balance that regulates glycemia). Its potential role must be distinguished from the general picture. This includes metabolic dysfunction (alteration of the processes that use energy), mild inflammation, and changes in the liver and adipose tissue. If the context changes, the weight of the data changes as well. The same result can have a different value in people with different metabolic profiles. The criterion is simple: first observe the context, then evaluate the ingredient. The following points help to understand this:

- The metabolic context changes the interpretation of studies on turmeric.
- The response to insulin guides research on sugar utilization.
- A single ingredient cannot explain such a complex dysfunction.

This section explains why your starting point matters just as much as the ingredient being studied. The three steps separate the context, the biological data, and the interpretation of the evidence.

6.1 Why metabolic context changes the interpretation of studies on turmeric

Studies do not show the same effects in all people. Your initial metabolic profile can change the observed response and how it is interpreted. Those who struggle with glucose management or have different insulin levels may react differently. The same applies in the presence of other metabolic signals. Understanding insulin resistance therefore helps to link each data point to the biological starting point. Every result only becomes meaningful when compared with well-defined initial metabolic conditions. In this way, the general picture, individual factors, and the interpretation of the evidence are kept distinct. Data must be read based on the initial profile, as responses are not uniform. Therefore, it is not easy to extend a result to everyone. This distinction prevents attributing differences to turmeric that actually depend on individual profiles. This limitation weighs even more heavily when an ingredient is isolated from a complex system. The next step concerns the insulin response in metabolism studies[10].

6.2 What role does the ability to respond well to insulin play in metabolism research

The ability of tissues to respond well to insulin indicates how effectively they react to the signal of this hormone. In metabolism studies, it helps determine whether an intervention regulates blood glucose. It also allows for distinguishing this effect from changes in less direct data. Regarding turmeric, researchers evaluate potential links to an already altered system. It is therefore necessary to observe this data alongside other signals that describe the same metabolic balance. However, such a hypothesis alone does not demonstrate a clinical effect in humans. This responsiveness, however, cannot be described by a single value or measure. In the same person, it can vary over time based on diet and physical activity. Low-grade inflammation, the liver, and adipose tissue also play a role. The data therefore offers a key to interpretation, but it is not final proof. Only this joint reading clarifies what each change detected in the study truly measures. Research evaluates how sugar utilization changes under various conditions, without providing answers that are valid for everyone.

6.3 Why we should not simplify the effect of a single ingredient

A single ingredient cannot necessarily have the same effect on every person's metabolism. Glycemic homeostasis (the balance that keeps glycemia stable) depends on many tissues and hormones. It also includes a network of responses involving the liver and adipose tissue. Evaluating a single effect therefore requires considering the other parts of the network as well. For this reason, turmeric must be studied within the context of an already existing metabolic dysfunction. It cannot be viewed as a solution that acts

on its own. Research shows genuine interest in these ingredients, but a methodological limitation remains. A single factor cannot explain a system made up of many parts acting together. Even when useful changes emerge, they should not be separated from the context in which they appear. Only in this way does an observed change maintain its significance without becoming a general conclusion. Their weight, in fact, depends on the initial condition and the duration of the study. This ingredient can therefore be part of the scientific discussion, but it does not replace a complete and cautious interpretation of the evidence.

7. QUADRO COMPARATIVO E SINTESI DELLE EVIDENZE

This summary collects the most representative sources from the bibliography. The comparison shows the evidence regarding the tissues involved, inflammation, measurement methods, and the general metabolic framework.

Sintesi delle evidenze scientifiche

Publication year	Area of study	What the studies indicate
2020	Tissue cross-talk	evidence indicates that muscle, liver, and adipose tissue communicate to control whole-body metabolism [10]
2022	Assessment of insulin response	studies suggest that HOMA-IR (an index based on glycemia and insulin) helps estimate insulin resistance in research and clinical practice [1]
2023	Adipose tissue and metabolism	studies suggest that adipose tissue regulates the body's energy and that its alterations impact metabolic diseases [2]
2024	Adipose tissue and liver	evidence indicates that altered adipose tissue promotes liver insulin resistance and metabolic disorder [5]
2025	Inflammation and insulin resistance	evidence indicates that the two processes are linked, but are not the same thing [8]
2025	Skeletal muscle	evidence indicates that muscle plays a central role in glucose uptake and the response to insulin [3]

8. SINTESI EDITORIALE E CONCLUSIONI

EDITORIAL INTERPRETATION OF THE EVIDENCE

Reading the studies collected in this article, I found a consistent picture: insulin resistance is described as a widespread process, not an isolated event. The data agree on the role of muscle, liver, and adipose tissue, which regulate glucose and energy in different ways. Recent reviews indicate to me that the dialogue between organs influences the response to insulin. The location of fat and low-grade inflammation also matter. From this, I have formed the conviction that metabolism changes due to the action of multiple signals and does not depend on a single biological switch.

I must note, however, that the bibliography examined here consists primarily of reviews and narrative syntheses. Experimental studies measuring the same outcomes are less common, and many abstracts do not report all numerical data. This limitation makes a direct comparison difficult for me. Furthermore, individual differences do not allow for setting thresholds that are valid for everyone, nor for giving the same weight to every tissue. I therefore consider the model theoretically solid, but it requires caution in

estimates and predictions.

FINAL CRITICAL NOTE

The general framework I have outlined is solid, but the available evidence does not all carry the same weight. Reviews help me clarify the mechanisms, but they primarily describe theoretical links and models. The boundary between useful adaptation and lasting damage remains less clear to me, and the weight of the various factors is not yet well defined. I therefore suggest caution: metabolism does not follow a single path. Furthermore, I have observed that the value of biological signals changes based on the individual and the period observed.

CONCLUSIONS AND EDITORIAL SUMMARY

From this analysis, I have concluded that insulin resistance indicates a change in the biological dialogue between insulin and tissues. It is neither a single label nor a simple laboratory anomaly. To assess its impact, I have observed that it is necessary to consider the muscle, liver, and adipose tissue together, as they manage the majority of glucose and energy. The picture, therefore, goes beyond glycemia and concerns homeostasis, that is, the internal balance of the body.The most recent studies collected here describe a dynamic, varied, and context-dependent process. For this reason, I believe that overly simplistic interpretations are of little use. What matters is the location of the fat, the inflammatory state, and the dialogue between tissues. This perspective, in my opinion, also helps in understanding future articles on metabolism and in studying the relationships between nutrients, cellular signals, and metabolic profile.

Aggiornato al 24/11/2026. Questo contenuto riflette una sintesi divulgativa delle evidenze scientifiche al momento della pubblicazione e non sostituisce il parere medico.

9. TRANSPARENCY AND SCIENTIFIC RESPONSIBILITY

Artificial Intelligence was used solely as a tool for editorial and documentary support. All content, interpretations, bibliographic citations and conclusions have been verified, reviewed and approved by the author, who assumes full scientific responsibility for them.

The manuscript was also subjected to independent peer review prior to publication.

To ensure the greatest transparency, the complete history of the article, including the dates of submission, revision, acceptance and publication, as well as information on the peer review process, is available on the dedicated Article History page.

10. RIFERIMENTI BIBLIOGRAFICI

[1] **HOMA-IR mean values in healthy individuals: a population-based study in iranian subjects**
Masoodian et al., 2022
Journal of Diabetes & Metabolic Disorders
DOI: https://doi.org/10.1007/s40200-022-01099-9

[2] **Insulin resistance in adipose tissue and metabolic diseases**
Imi et al., 2023
Diabetology International
DOI: https://doi.org/10.1007/s13340-022-00616-8

[3] **Unraveling Skeletal Muscle Insulin Resistance: Molecular Mechanisms and the Restorative Role of Exercise**
Whytock et al., 2025
Circulation Research
DOI: https://doi.org/10.1161/CIRCRESAHA.125.325532

[4] **Insulin resistance and adipose tissue interactions as the cornerstone of metabolic (dysfunction)-associated fatty liver disease pathogenesis**
Pal et al., 2023
World Journal of Gastroenterology
DOI: https://doi.org/10.3748/wjg.v29.i25.3999

[5] **The role of adipose tissue dysfunction in hepatic insulin resistance and T2D**
Sancar et al., 2024
Journal of Endocrinology
DOI: https://doi.org/10.1530/JOE-24-0115

[6] **Insulin resistance and steatosis in humans**
Capeau et al., 2008
Diabetes & Metabolism
DOI: https://doi.org/10.1016/S1262-3636(08)74600-7

[7] **Additive Interaction Between Insulin Resistance, Chronic Low-Grade Inflammation and Vitamin D Deficiency on the Risk of Type 2 Diabetes Mellitus: A Cohort Study**
Bennouar et al., 2024
Journal of the American Nutrition Association
DOI: https://doi.org/10.1080/27697061.2024.2352401

[8] **The close relationship between inflammation and insulin resistance: a comparative perspective from a new angle**
Wang et al., 2025
Acta Diabetologica
DOI: https://doi.org/10.1007/s00592-025-02596-y

[9] **Quantification of Adipose Tissue Insulin Sensitivity**
Søndergaard et al., 2016
Journal of Investigative Medicine
DOI: https://doi.org/10.1136/jim-2016-000098

[10] **Mechanisms of muscle insulin resistance and the cross talk with liver and adipose tissue**
Silva Rosa et al., 2020
Physiological Reports
DOI: https://doi.org/10.14814/phy2.14607

[11] **The Vicious Cycle of Type 2 Diabetes Mellitus and Skeletal Muscle Atrophy: Clinical, Biochemical, and Nutritional Bases**
Lopez-Pedrosa et al., 2024
Nutrients
DOI: https://doi.org/10.3390/nu16010172