LIFE Science Hub
JOURNAL OF SCIENCE
EXPANDING KNOWLEDGE ABOUT NUTRITION AND HEALTH

ORIGINAL RESEARCH
ISSN 2845-7501

Insulin stimulation of SREBP-1c processing in transgenic rat hepatocytes requires p70 S6-kinase - Review of the evidence

D.ssa D. Iodice[1]

1 Biologist, Italy

1. ABSTRACT

The analyzed study evaluated how insulin activates SREBP-1c, a transcription factor that turns on genes involved in the synthesis of fatty acids and triglycerides in the liver. In freshly isolated hepatocytes from transgenic rats, the authors separated two levels of regulation: the production of SREBP-1c messenger RNA and its proteolytic processing, which is necessary to release the form capable of entering the nucleus. The results indicate that both depend on mTORC1, while processing also requires p70 S6-kinase. The number of experiments, dosages, and quantitative data are not reported in the available material, which consists only of the abstract.In section 5.3 of "The Choice to Live in Health or Illness" by Dr. A. Colonnese and R. Panzironi, the publication contributes to the passage stating that insulin exerts a dual control—transcriptional and post-translational—on SREBP-1c. Independent research confirms that insulin increases SREBP-1c expression via mTORC1 and that it can also increase its proteolytic maturation. Other evidence shows that SREBP links AKT/mTORC1 signals to lipid synthesis and cell growth. The confirmation is primarily mechanistic, as it is derived from cells and animal models.

ONLINE ARTICLE

ARTICLE INFORMATION

Published: November 23, 2026
Updated: November 23, 2026

DOI 10.5281/zenodo.24470333

MISSION

LIFE Science Hub – Journal of Science was founded with the aim of making scientific knowledge accessible without giving up methodological rigour, quality of sources and authoritativeness of content. The journal promotes the dissemination of knowledge in the health, nutrition, medical and longevity sciences, building a bridge between the world of research and society.

Through articles, reviews, in-depth analyses and expert contributions, the Journal champions evidence-based medicine and science, fostering a correct interpretation of scientific data and countering misinformation. Every piece of content is developed with the aim of translating the complexity of research into understandable, useful and applicable information, while keeping intact the principles of accuracy, transparency and scientific independence.

Free from political, industrial and commercial influence, the journal stands as an authoritative and independent editorial space, dedicated to promoting scientific culture, professional development and the growth of health awareness among the public.

2. ANALYZED STUDY

The research question was whether insulin only controls the amount of SREBP-1c produced by the cell or also its final activation. SREBP-1c is initially an inactive protein embedded in the membrane of the endoplasmic reticulum, an internal network within the cell. To become operational, it must be transferred and cleaved by enzymes; the released fragment can then reach the nucleus and increase the expression of lipogenesis genes, i.e., lipid production.The study is relevant to section 5.3 of "The Choice to Live in Health or Illness" by Dr. A. Colonnese and R. Panzironi because it precisely documents the second level of control described in the text: the post-translational regulation of SREBP-1c by insulin. In particular, the publication distinguishes the pathway that increases SREBP-1c messenger RNA from the pathway that promotes its proteolytic processing.The research can demonstrate a causal relationship in the experimental model used, because it compares cells treated with insulin and cells exposed to inhibitors of the molecular pathways considered. However, it cannot establish that the same mechanism has identical intensity in the human liver, nor prove that chronic hyperinsulinemia directly causes a specific disease in humans.

3. MATERIALS AND METHODS OF THE ORIGINAL STUDY

The study used a transgenic rat line engineered to produce a human form of SREBP-1c in the liver, which could be experimentally identified using a protein tag. Protein expression was driven by an apolipoprotein E promoter, chosen to keep SREBP-1c production separate from the increase in its transcription induced by insulin.The authors isolated fresh hepatocytes, i.e., liver cells, from the transgenic rats. The cells were exposed to insulin and to pharmacological interventions targeting mTORC1, the protein complex that integrates nutrient and growth signals, and p70 S6-kinase, a protein activated downstream of mTORC1. The effect of glucagon, a hormone that counteracts insulin action in many metabolic conditions, was also evaluated.Endpoints included SREBP-1c messenger RNA and the processing of the precursor protein into its active form. The number of animals, the number of independent cultures, inclusion and exclusion criteria, inhibitor concentrations, treatment duration, detailed analytical techniques, statistical analyses, and significance values are not reported in the available material.

4. RESULTS

The main finding is that insulin increased SREBP-1c processing in transgenic rat hepatocytes. In simple terms, the hormone did not just increase the starting material of the protein, but also promoted its transition into the form that can enter the nucleus and activate lipogenic genes.According to the abstract, mTORC1 inhibition blocked both the insulin-induced increase in SREBP-1c messenger RNA and its processing. This indicates that mTORC1 is required for both levels of regulation in the studied model. Conversely, inhibition of p70 S6-kinase blocked proteolytic processing but not the induction of messenger RNA. Therefore, after mTORC1, the two responses follow at least partially different pathways.Glucagon blocked both of the insulin effects reported in the abstract: the increase in messenger RNA and the processing of SREBP-1c. No absolute values, percentage variations, P-values, or confidence intervals are reported. Therefore, the result allows for the identification of the direction of the effects and the functional relationship between the steps, but not for the estimation of their quantitative magnitude.

5. INTERPRETATION

For a liver cell, increasing SREBP-1c means enhancing the ability to transform energy availability signals into the production of lipid components. The study indicates that insulin acts on this protein in a dual manner: it increases the production of its messenger RNA and promotes the cleavage of the precursor protein into its active form.The segment specifically documented by the publication subject to the Evidence Review is: insulin mTORC1 p70 S6-kinase increased processing of SREBP-1c. The p70 S6-kinase appears necessary for the post-translational branch in the transgenic hepatocyte model, while the increase in SREBP-1c messenger RNA depends on mTORC1 without requiring this kinase.This distinction supports the book's narrative that SREBP-1c is particularly sensitive to a persistent insulin signal because it can be regulated both before and after protein synthesis. However, the study does not measure chronic insulin

exposure in humans and does not directly demonstrate the inflammatory or clinical consequences of increased lipogenesis.

6. INDEPENDENT CONFIRMING EVIDENCE

Initial confirmation concerns the transcriptional branch. In rat hepatocytes, insulin increased SREBP-1c messenger RNA by 25-fold and reduced the messenger RNA of PEPCK, an enzyme involved in hepatic glucose production, by 95%. PI3K and AKT inhibitors blocked both responses, while rapamycin selectively blocked the induction of SREBP-1c without preventing the reduction of PEPCK. The same selective effect of rapamycin was observed in the livers of rats and mice after fasting and refeeding. This study confirms that mTORC1 is necessary for the lipogenic component of the insulin response, but it does not establish the role of proteolytic processing on its own [1].A second study directly examined the post-translational step in primary rat hepatocytes. Insulin increased the nuclear accumulation of the active form of SREBP-1c by 2.62 ± 0.21-fold and more than doubled the phosphorylation of the precursor protein. PI3K and AKT inhibitors prevented both phosphorylation and processing. The results therefore link insulin to the migration of SREBP-1c from the endoplasmic reticulum to the Golgi and its subsequent proteolytic cleavage. This work does not identify mTORC1 or p70 S6-kinase as mandatory steps, but it independently confirms that insulin regulates the maturation of SREBP-1c in addition to its expression [2].A third study expands the functional significance of the chain. In immortalized human epithelial cells, AKT activation increased lipid synthesis and the accumulation of mature SREBP in the nucleus; rapamycin blocked these effects. Silencing SREBP reduced AKT-induced lipogenesis and attenuated the increase in cell size. In Drosophila, silencing the SREBP homolog reduced cell and organ size and prevented the growth increase induced by PI3K. This evidence confirms that SREBP is a functional component, not just an indicator, of the anabolic program activated by AKT and mTORC1. The limitation is that the model does not specifically concern the liver or insulin in a physiological context [3].

7. CONFLICTING EVIDENCE

No directly conflicting evidence of sufficient relevance has been identified; this does not prove the absence of contrary results.The main interpretative caution concerns the dependence on the experimental context. Available studies show that mTORC1 is necessary for specific insulin responses of SREBP-1c in hepatocytes and rodents, but they do not allow it to be considered the sole regulator of lipogenesis. Nutrients, glucose, other transcription factors, and hormonal signals can modulate the same network. Furthermore, cell culture studies verify causal mechanisms under controlled conditions, but they do not fully reproduce the complexity of chronic hyperinsulinemia in the human body.

8. QUADRO COMPARATIVO E SINTESI DELLE EVIDENZE

The table distinguishes the study that is the subject of the Evidence Review from the independent publications used to confirm the individual steps of the narrative.

Quadro comparativo

Reference	Study type	Population or model	Main result	Direction of evidence	Strength of evidence
Original study	Experimental cellular	Fresh hepatocytes from transgenic rats	Insulin increases SREBP-1c processing; mTORC1 is necessary and p70 S6-kinase is required for the post-translational branch.	direct confirmation	moderate
[1]	Experimental cellular and animal	Rat hepatocytes; rat and mouse liver after fasting-refeeding	Insulin increases SREBP-1c messenger RNA via PI3K/AKT/mTORC1; rapamycin blocks this increase.	direct confirmation	moderate
[2]	Experimental cellular	Primary rat hepatocytes and rat hepatoma cells	Insulin increases phosphorylation, transport, and proteolytic cleavage of SREBP-1c; PI3K and AKT are necessary.	direct confirmation	moderate
[3]	Experimental cellular and animal	Immortalized human epithelial cells and Drosophila	mTORC1 and SREBP are necessary for lipogenesis and part of the cell growth induced by AKT.	biologically consistent evidence	moderate

9. SINTESI EDITORIALE E CONCLUSIONI

CONCLUSIONS

The study analyzed precisely supports the concept that insulin controls SREBP-1c not only by increasing its expression but also by promoting its maturation into the active form. The role of mTORC1 is well-supported in rodent hepatic models, while the dependence of the processing on p70 S6-kinase is mechanistically supported in the transgenic model examined.Independent evidence strengthens the link between insulin signaling, PI3K/AKT, mTORC1, SREBP-1c activation, and lipogenesis. However, most of the data are derived from cells and animals. Extending these results to human physiology and the consequences of chronic hyperinsulinemia requires clinical validation and studies in human tissues.

Aggiornato al 24/11/2026. Questo contenuto riflette una sintesi divulgativa delle evidenze scientifiche al momento della pubblicazione e non sostituisce il parere medico.

10. TRANSPARENCY AND SCIENTIFIC RESPONSIBILITY

Artificial Intelligence was used solely as a tool for editorial and documentary support. All content, interpretations, bibliographic citations and conclusions have been verified, reviewed and approved by the author, who assumes full scientific responsibility for them.

The manuscript was also subjected to independent peer

review prior to publication.
To ensure the greatest transparency, the complete history of the article, including the dates of submission, revision, acceptance and publication, as well as information on the peer review process, is available on the dedicated Article History page.

11. RIFERIMENTI BIBLIOGRAFICI

[1] **Bifurcation of insulin signaling pathway in rat liver: mTORC1 required for stimulation of lipogenesis, but not inhibition of gluconeogenesis**
Li et al., 2010
Proceedings of the National Academy of Sciences
DOI: https://doi.org/10.1073/pnas.0914798107

[2] **Insulin Enhances Post-translational Processing of Nascent SREBP-1c by Promoting Its Phosphorylation and Association with COPII Vesicles**
Yellaturu et al., 2009
Journal of Biological Chemistry
DOI: https://doi.org/10.1074/jbc.M805746200

[3] **SREBP Activity Is Regulated by mTORC1 and Contributes to Akt-Dependent Cell Growth**
Porstmann et al., 2008
Cell Metabolism
DOI: https://doi.org/10.1016/j.cmet.2008.07.007