LIFE Science Hub
JOURNAL OF SCIENCE
EXPANDING KNOWLEDGE ABOUT NUTRITION AND HEALTH

ORIGINAL RESEARCH
ISSN 2845-7501

Rottlerin enhances IL-1b-induced COX-2 expression through sustained activation of p38 MAPK in human breast carcinoma MDA-MB-231 cells - Evidence review

Dott. A. Colonnese[1]

1 Nutritional Biologist, Italy

1. ABSTRACT

The analyzed study examined whether rottlerin enhanced the increase of cyclooxygenase-2 (COX-2), an enzyme involved in prostaglandin production, induced by interleukin-1b (IL-1b), a molecule that promotes the inflammatory response. The research was conducted in a single human breast carcinoma cell line, MDA-MB-231. The combination of the two stimuli increased messenger RNA and the COX-2 protein; the authors attributed part of the effect to the increased stability of the messenger RNA and the prolonged activation of the p38 MAPK protein.In paragraph 5.5 of “The Choice to Live in Health or Illness” by Dr. A. Colonnese and R. Panzironi, the publication contributes to the narrative that MAPK activation can increase COX-2 and support a lipid inflammatory response. However, the original study documents p38 MAPK, not specifically the ERK branch. Independent studies confirm that p38 can stabilize COX-2 RNA and promote its expression; other data show that the relative contribution of p38 and ERK varies with cell type and stimulus. The confirmation is therefore primarily mechanistic and cellular.

ONLINE ARTICLE

ARTICLE INFORMATION

Published: November 23, 2026
Updated: November 23, 2026

DOI 10.5281/zenodo.24459202

MISSION

LIFE Science Hub – Journal of Science was founded with the aim of making scientific knowledge accessible without giving up methodological rigour, quality of sources and authoritativeness of content. The journal promotes the dissemination of knowledge in the health, nutrition, medical and longevity sciences, building a bridge between the world of research and society.

Through articles, reviews, in-depth analyses and expert contributions, the Journal champions evidence-based medicine and science, fostering a correct interpretation of scientific data and countering misinformation. Every piece of content is developed with the aim of translating the complexity of research into understandable, useful and applicable information, while keeping intact the principles of accuracy, transparency and scientific independence.

Free from political, industrial and commercial influence, the journal stands as an authoritative and independent editorial space, dedicated to promoting scientific culture, professional development and the growth of health awareness among the public.

2. ANALYZED STUDY

The research question was whether rottlerin could amplify the induction of COX-2 caused by IL-1b and, if so, through which intracellular signal. COX-2 is an inducible enzyme: when it increases, the cell acquires a greater capacity to convert arachidonic acid into prostanoids, lipid molecules that include the precursors of prostaglandin E2 (PGE2).The hypothesis evaluated was that persistent activation of p38 MAPK, a protein that transmits stress and inflammation signals, would make COX-2 messenger RNA more durable. In simple terms, rather than necessarily producing much more new RNA, the cell could degrade it more slowly and thus maintain the instructions needed to build the enzyme for longer.The relevance to paragraph 5.5 consists of the segment of the chain that connects a MAPK pathway to the increase in COX-2. The analyzed study supports this segment in a human tumor cell model and in the presence of inflammatory stimuli. It does not evaluate insulin, does not directly measure the availability of arachidonic acid, and does not demonstrate that the mechanism occurs in the tissues of healthy or sick people. Furthermore, the documented MAPK is p38, while the book's paragraph expressly cites MAPK-ERK.

3. MATERIALS AND METHODS OF THE ORIGINAL STUDY

The study was conducted on the MDA-MB-231 cell line, derived from human breast carcinoma. The cells were treated with rottlerin, IL-1b, or a combination of both. The authors evaluated COX-2 at both the protein and messenger RNA levels.To clarify the potential mechanism, the stability of COX-2 messenger RNA was assessed. The activation of p38 MAPK was also examined. The authors used SB203580, a pharmacological inhibitor of p38 MAPK, and temporarily introduced an inactive form of p38 MAPK into the cells. Direct experiments on protein kinase C delta (PKCδ) were also performed using small interfering RNA (siRNA) and a dominant-negative form of the protein, which is designed to hinder its function.Finally, other pro-inflammatory stimuli were tested: tumor necrosis factor-alpha, phorbol ester, and bacterial lipopolysaccharide. The number of experiments, concentrations, duration of treatments, statistical criteria, P-values, and full technical details are not reported in the available material.

4. RESULTS

Rottlerin increased IL-1b-induced COX-2 expression at both the messenger RNA and protein levels. This result means that, in these laboratory-cultured tumor cells, the combination of the two stimuli produced more molecular instructions for COX-2 and more detectable enzyme compared to the stimulation with IL-1b considered by the authors.The authors observed that the increase depended, at least in part, on greater stability of the COX-2 messenger RNA. More stable RNA persists longer in the cell and can be translated into protein multiple times. In parallel, rottlerin and IL-1b induced sustained activation of p38 MAPK. Pharmacological inhibition of p38 with SB203580 and the expression of an inactive form of p38 reduced the increase in COX-2 induced by the combination.The reduction of PKCδ via siRNA, as well as the expression of its dominant-negative form, did not abolish the increase in COX-2. This result makes PKCδ less plausible as a necessary mediator of the effect observed under the experimental conditions considered. Rottlerin also enhanced the induction of COX-2 caused by TNF-a, phorbol ester, and lipopolysaccharide. The abstract does not report absolute values, percentages, confidence intervals, or P-values; therefore, such data cannot be described beyond what the authors have stated.

5. INTERPRETATION

The most direct biological significance is that an inflammatory signal can not only initiate COX-2 expression but also prolong it by reducing the degradation of its messenger RNA. In a cell, this can make the availability of the enzyme involved in prostanoid production more persistent. The analyzed study identifies p38 MAPK as a functional element of this regulation: when p38 was pharmacologically blocked or rendered inactive, the increase in COX-2 was reduced.This result is consistent with independent studies in which the inhibition of p38 destabilized IL-1-induced COX-2 RNA and reduced PGE2 production [1]. A separate experimental model also indicated that the p38 cascade can stabilize regulatory sequences present in the terminal region of

COX-2 RNA through MAPKAPK-2, a protein activated by p38 [2].The crucial interpretative limitation concerns the terminology in the book's paragraph. p38 and ERK both belong to the MAPK family, but they are distinct branches. The publication subject to the Evidence Review documents p38 MAPK and does not demonstrate that sustained ERK activation stabilizes COX-2 RNA in its model. The general MAPK COX-2 link is therefore supported; the specific attribution of the effect described in this publication to MAPK-ERK is not faithful to the data from the original study.

6. INDEPENDENT SUPPORTING EVIDENCE

The first independent confirmation concerns the step closest to the analyzed publication: IL-1 p38 MAPK COX-2 RNA stability PGE2. Ridley and colleagues studied HeLa cells, a human cell line, stimulated with IL-1. The p38 inhibitor SB203580 reduced the induction of COX-2 RNA and protein and reduced the release of PGE2, without significantly decreasing COX-2 transcription. When the RNA was already being produced, the inhibitor instead promoted its rapid destabilization. This data directly confirms, albeit only at the cellular level, that p38 can maintain the availability of COX-2 RNA following an inflammatory stimulus [1].A second study investigated the mechanism in greater depth. Lasa and colleagues used HeLa-TO cells and an experimental system in which RNA was produced in a controllable manner. Activation of the p38 cascade stabilized a reporter RNA containing the 3 untranslated region of COX-2, which is a part of the RNA that helps determine its lifespan and degradation. The effect required MAPKAPK-2, a protein that acts downstream of p38. This research does not concern breast cancer or insulin, but it makes the interpretation that p38 can prolong the life of COX-2 RNA biologically more robust [2].Taken together, these independent pieces of evidence support a convergent cellular chain: an inflammatory stimulus activates p38, p38 makes COX-2 RNA more stable, and this promotes greater availability of the enzyme. The level of confirmation remains mechanistic: the models are cell cultures and not clinical or observational studies in humans.

7. CONFLICTING EVIDENCE

No directly conflicting evidence sufficiently relevant to the fact that p38 may contribute to COX-2 RNA stability has been identified. However, there is significant data that limits the generalizability of the specific pathway involved.In human A549 lung epithelial cells, the increase in COX-2 and the release of PGE2 induced by phorbol ester were reduced by the inhibition of the Ras/Raf/ERK and NF-kB pathways, whereas the inhibition of p38 did not produce the same reduction [3]. This does not contradict studies focused on IL-1b and p38; rather, it shows that COX-2 regulation depends on the cell type and the stimulus. In some contexts, p38 prevails, while in others, ERK does; furthermore, multiple pathways can converge on the same gene.This context-dependence is particularly relevant for the analyzed paragraph. It is incorrect to treat MAPK-ERK and p38 MAPK as if they were interchangeable. Both can participate in the induction of COX-2, but their relative weight, the mechanism involved, and the effect on RNA stability must be verified in the specific model.

8. QUADRO COMPARATIVO E SINTESI DELLE EVIDENZE

The table distinguishes the publication subject to the review from independent evidence and indicates the level at which each result supports the narrative of the paragraph.

Quadro comparativo

Reference	Study type	Population or model	Main result	Direction of evidence	Strength of evidence
Original study	Experimental cellular	Human breast carcinoma cells MDA-MB-231	Rottlerin and IL-1b increase COX-2; the effect involves RNA stability and sustained activation of p38 MAPK.	biologically consistent evidence	preliminary
[1]	Experimental cellular	Human HeLa cells stimulated with IL-1	p38 inhibition reduces COX-2 and PGE2 and destabilizes COX-2 RNA.	direct confirmation	moderate
[2]	Mechanistic cellular	HeLa-TO cells with inducible reporter RNA	The p38 cascade stabilizes COX-2 RNA regulatory sequences via MAPKAPK-2.	direct confirmation	moderate
[3]	Experimental cellular	Human lung epithelial cells A549	Induction of COX-2 and PGE2 by phorbol ester depends on ERK, whereas p38 is not required in that model.	discordant evidence	moderate

9. SINTESI EDITORIALE E CONCLUSIONI

CONCLUSIONS

The study analyzed maintains that, in MDA-MB-231 cells, rottlerin enhances the induction of COX-2 by IL-1b through a mechanism that includes sustained activation of p38 MAPK and increased stability of COX-2 messenger RNA. This segment of the narrative is confirmed by independent evidence at the cellular and molecular level.The most accurate formulation is that p38 MAPK can contribute to prolonging COX-2 expression in specific inflammatory contexts. It is not demonstrated by the study that insulin produces the same effect, nor that the mechanism necessarily involves ERK, nor that it determines clinical consequences. The narrative is well-supported on a mechanistic level, but requires validation in physiological models and in humans.

Aggiornato al 24/11/2026. Questo contenuto riflette una sintesi divulgativa delle evidenze scientifiche al momento della pubblicazione e non sostituisce il parere medico.

10. TRANSPARENCY AND SCIENTIFIC RESPONSIBILITY

Artificial Intelligence was used solely as a tool for editorial and documentary support. All content, interpretations, bibliographic citations and conclusions have been verified, reviewed and approved by the author, who assumes full scientific responsibility for them.

The manuscript was also subjected to independent peer review prior to publication.

To ensure the greatest transparency, the complete history of the article, including the dates of submission, revision, acceptance and publication, as well as information on the peer review process, is available on the dedicated Article History page.

11. RIFERIMENTI BIBLIOGRAFICI

[1] **A p38 MAP kinase inhibitor regulates stability of interleukin 1 induced cyclooxygenase 2 mRNA**
Ridley et al., 1998
FEBS Letters
DOI: https://doi.org/10.1016/S0014-5793(98)01342-8

[2] **Regulation of Cyclooxygenase 2 mRNA Stability by the Mitogen-Activated Protein Kinase p38 Signaling Cascade**
Lasa et al., 2000
Molecular and Cellular Biology
DOI: https://doi.org/10.1128/MCB.20.12.4265-4274.2000

[3] **Phorbol 12-myristate 13-acetate upregulates cyclooxygenase-2 expression in human pulmonary epithelial cells via Ras, Raf-1, ERK, and NF-kB, but not p38 MAPK, pathways**
Chang et al., 2005
Cellular Signalling
DOI: https://doi.org/10.1016/j.cellsig.2004.07.008