LIFE Science Hub
JOURNAL OF SCIENCE
EXPANDING KNOWLEDGE ABOUT NUTRITION AND HEALTH

ORIGINAL RESEARCH
ISSN 2845-7501

Sleep loss: a new risk factor for insulin resistance and type 2 diabetes - Evidence review

D.ssa D. Iodice[1]

1 Biologist, Italy

1. ABSTRACT

The analyzed study is a narrative synthesis examining how chronic or recurrent sleep loss can influence glucose metabolism. The available abstract describes experiments conducted on healthy young adults, in which partial sleep restriction was associated with reduced glucose tolerance and lower insulin sensitivity. It also describes neuroendocrine alterations, including changes in leptin and ghrelin, and cites epidemiological evidence on the association between short sleep, obesity, and type 2 diabetes.In section 4.15 of “The Choice to Live in Health or Illness” by Dr. A. Colonnese and R. Panzironi, the publication contributes to the narrative that reduced sleep duration can promote insulin resistance and alterations in the stress axis. Independent literature consistently confirms that experimental sleep restriction reduces insulin sensitivity. The specific link with an increase in cortisol, however, is less uniform: some studies suggest it may contribute, but they do not isolate cortisol as the sole cause of the metabolic effect.

ONLINE ARTICLE

ARTICLE INFORMATION

Published: November 20, 2026
Updated: November 20, 2026

DOI 10.5281/zenodo.24403569

MISSION

LIFE Science Hub – Journal of Science was founded with the aim of making scientific knowledge accessible without giving up methodological rigour, quality of sources and authoritativeness of content. The journal promotes the dissemination of knowledge in the health, nutrition, medical and longevity sciences, building a bridge between the world of research and society.

Through articles, reviews, in-depth analyses and expert contributions, the Journal champions evidence-based medicine and science, fostering a correct interpretation of scientific data and countering misinformation. Every piece of content is developed with the aim of translating the complexity of research into understandable, useful and applicable information, while keeping intact the principles of accuracy, transparency and scientific independence.

Free from political, industrial and commercial influence, the journal stands as an authoritative and independent editorial space, dedicated to promoting scientific culture, professional development and the growth of health awareness among the public.

2. ANALYZED STUDY

The question addressed is whether habitual sleep loss, whether due to behavioral choices or sleep disorders, can contribute to insulin resistance and type 2 diabetes. The biological hypothesis is intuitive: if sleep is insufficient, the body may respond less effectively to insulin, the hormone that facilitates the entry of glucose into cells.The publication is relevant to sections 4.14 and 4.15 of "The Choice to Live in Health or Illness" by Dr. A. Colonnese and R. Panzironi because it describes the segment of the chain that links sleep loss to reduced insulin sensitivity. In the available abstract, the authors also report neuroendocrine changes and discuss the possible role of sleep fragmentation and hypoxia in metabolic disorders.However, the research covered in the article is not presented as a single observational study with participants, a six-hour sleep threshold, morning cortisol levels, and statistical corrections for body mass index, age, and diet. These details are not reported in the available material. The strongest correspondence therefore concerns the general statement: sleep restriction can worsen glucose regulation and promote insulin resistance; it does not directly demonstrate all the clinical and hormonal details attributed to the reference in the book.

3. MATERIALS AND METHODS OF THE ORIGINAL STUDY

The available material consists only of the abstract. The publication appears to be a narrative review: it summarizes the results of laboratory studies on healthy young adults and epidemiological studies, without presenting a unique experimental protocol, an original cohort, or a new statistical analysis in the available text.The overall sample size, inclusion and exclusion criteria, observation periods, analytical techniques, predefined endpoints, and statistical procedures are not reported in the available material. The abstract reports that, in the experiments cited by the authors, healthy participants were subjected to partial and recurrent sleep restriction; however, it does not specify the duration, number of nights, or characteristics of the individual protocols.The indicators discussed include glucose tolerance, insulin sensitivity, leptin, and ghrelin. Leptin is a hormone that contributes to the feeling of satiety, while ghrelin promotes the sensation of hunger. The available text does not allow for the determination of which methods were used in the individual studies to measure these indicators.

4. RESULTS

The key finding reported in the abstract is that recurrent partial sleep restriction, in experiments on healthy young adults, was accompanied by a reduction in glucose tolerance and insulin sensitivity. In simple terms, after insufficient sleep, the body managed glucose less effectively and required greater insulin action to achieve a comparable metabolic response.The abstract also reports a decrease in leptin and an increase in ghrelin. This direction of effect is consistent with increased hunger and appetite, which the authors indicate as possible factors promoting overeating and weight gain. However, no absolute values, percentage changes, confidence intervals, or P-values are reported.The authors also describe growing epidemiological evidence of an association between short sleep duration and the risk of obesity or diabetes. However, an observational association does not prove that short sleep alone causes diabetes: body weight, shift work, sleep apnea, diet, physical activity, and stress conditions can all contribute to the risk simultaneously. The abstract does not provide specific data on subjects who slept less than six hours, nor direct data on morning cortisol levels.

5. INTERPRETATION

The biological significance is that sleep does not only affect perceived rest. It can influence how the liver, muscles, and adipose tissue respond to insulin. If these tissues become less sensitive to the hormone, the body must produce or use more insulin to keep glucose within appropriate levels.The segment supported by the publication is: sleep reduction or loss worsening of insulin sensitivity higher probability of impaired glucose regulation. The hypothesis that cortisol participates in this pathway is physiologically plausible, because cortisol is a glucocorticoid involved in the stress response and can counteract some effects of insulin. However, the abstract does not provide direct experimental proof that an increase in cortisol is the sole or primary cause of insulin resistance induced by sleep loss.The wording of paragraph 4.15, which states that the reference documented an independent

association between sleep of less than six hours, higher morning cortisol, and a higher incidence of insulin resistance, should therefore be interpreted with caution. The study analyzed supports the general sleep-insulin resistance framework, but the available material does not directly document that specific observational analysis.

6. INDEPENDENT SUPPORTING EVIDENCE

Particularly direct experimental confirmation comes from a crossover study on nine healthy adults, five men and four women. Each participant was evaluated after a normal night's sleep and after a night limited to four hours of sleep. The restriction reduced the rate of glucose disposal in peripheral tissues and increased endogenous glucose production during a hyperinsulinemic-euglycemic clamp, a procedure that precisely measures the response to insulin. To maintain stable glycemia, approximately 25% less infused glucose was required after restricted sleep, a sign of lower insulin sensitivity. The study directly confirms the link between restricted sleep and insulin resistance, but the very small sample size and the single episode of restriction limit its generalizability [1].A systematic review with meta-analysis of randomized trials subsequently integrated 35 articles, of which 21 studies on sleep restriction were included in the specific meta-analysis. Sleep restriction reduced insulin sensitivity when assessed with glucose tolerance tests or HOMA-IR, an index calculated from fasting glycemia and insulin. Whole-body insulin sensitivity measured with a clamp was also found to be reduced. This evidence reinforces the consistency of the link described in the book, while not proving that every person with short sleep duration will develop type 2 diabetes [2].The possible hormonal involvement was explored more specifically in a randomized, double-blind, crossover study on 34 healthy young men subjected to four nights with four hours of sleep per night. Without hormonal intervention, the restriction increased insulinemia, glycemia, and insulin resistance. When the researchers simultaneously controlled exposure to cortisol and testosterone, the development of insulin resistance and hyperinsulinemia was attenuated by approximately 50%. The result is consistent with a role for the stress-related hormonal environment, but it does not separate the effect of cortisol from that of testosterone and does not demonstrate a single mechanism [3].

7. CONFLICTING EVIDENCE

No directly conflicting evidence sufficiently relevant to the general finding that experimental sleep restriction can reduce insulin sensitivity has been identified; this does not prove the absence of contrary results.However, there are findings that warrant caution regarding the specific role of cortisol. In the study on a single night of reduced sleep, insulin resistance increased during the clamp, but morning basal cortisol concentrations were not altered. This indicates that the reduced insulin response can occur even without a detectable increase in basal morning cortisol [1].Furthermore, the meta-analysis of randomized studies found an unfavorable effect of sleep duration restriction, circadian misalignment, and slow-wave sleep suppression; however, it did not find an overall effect of sleep fragmentation on insulin sensitivity. Peripheral insulin sensitivity was also not significantly reduced in the available pooled analysis. These differences suggest that duration, quality, sleep timing, intensity of the manipulation, and measurement method may modify the observed effect [2].The hormonal study attenuated, but did not eliminate, insulin resistance while keeping both cortisol and testosterone stable. Therefore, the data support a contribution from the hormonal profile, rather than the conclusion that cortisol alone explains the entire relationship between insufficient sleep and insulin resistance [3].

8. QUADRO COMPARATIVO E SINTESI DELLE EVIDENZE

The table distinguishes the publication being reviewed from the independent evidence used to verify the book's narrative.

Quadro comparativo

Reference	Study type	Population or model	Main result	Direction of evidence	Strength of evidence
Original study	Narrative review, material limited to the abstract	Previous studies on healthy young adults and epidemiological data discussed by the authors	Reports reduced glucose tolerance and insulin sensitivity following recurrent sleep restriction	biologically consistent evidence	moderate
[1]	Crossover experimental study	9 healthy adults; one normal night compared with one night of four hours of sleep	Reduction in hepatic and peripheral insulin sensitivity; approximately 25% less glucose infused during clamp	direct confirmation	moderate
[2]	Systematic review and meta-analysis of RCTs	35 eligible articles; 21 sleep restriction studies in the meta-analysis	Sleep restriction reduces insulin sensitivity; effects not uniform for fragmentation and peripheral sensitivity	direct confirmation	high
[3]	Double-blind crossover RCT	34 healthy young men; four nights with four hours of sleep	Combined control of cortisol and testosterone attenuates approximately 50% of restriction-induced insulin resistance and hyperinsulinemia	indirect confirmation	moderate

9. SINTESI EDITORIALE E CONCLUSIONI

CONCLUSIONS

Sleep loss or restriction is well-supported as a factor capable of reducing insulin sensitivity, at least temporarily, under controlled experimental conditions. The evidence is particularly consistent regarding insufficient sleep duration, while it is less uniform concerning sleep fragmentation and the individual biological mechanisms involved.The link with cortisol is biologically plausible and supported by experimental data on hormonal balance, but it requires further validation to distinguish the contribution of cortisol from that of testosterone, circadian rhythms, the autonomic nervous system, diet, and other factors. The original publication therefore supports the core of the narrative, but not all the specific details reported in the book's paragraph.

Aggiornato al 24/11/2026. Questo contenuto riflette una sintesi divulgativa delle evidenze scientifiche al momento della pubblicazione e non sostituisce il parere medico.

10. TRANSPARENCY AND SCIENTIFIC RESPONSIBILITY

Artificial Intelligence was used solely as a tool for editorial and documentary support. All content, interpretations, bibliographic citations and conclusions have been verified, reviewed and approved by the author, who assumes full scientific responsibility for them.

The manuscript was also subjected to independent peer review prior to publication.

To ensure the greatest transparency, the complete history of the article, including the dates of submission, revision, acceptance and publication, as well as information on the peer review process, is available on the dedicated Article History page.

11. RIFERIMENTI BIBLIOGRAFICI

[1] **A Single Night of Partial Sleep Deprivation Induces Insulin Resistance in Multiple Metabolic Pathways in Healthy Subjects**
Donga et al., 2010
The Journal of Clinical Endocrinology & Metabolism
DOI: https://doi.org/10.1210/jc.2009-2430

[2] **Effects of sleep manipulation on markers of insulin sensitivity: A systematic review and meta-analysis of randomized controlled trials**
Sondrup et al., 2022
Sleep Medicine Reviews
DOI: https://doi.org/10.1016/j.smrv.2022.101594

[3] **Clamping Cortisol and Testosterone Mitigates the Development of Insulin Resistance during Sleep Restriction in Men**
Liu et al., 2021
The Journal of Clinical Endocrinology & Metabolism
DOI: https://doi.org/10.1210/clinem/dgab375